


Trauma comes in many forms.
Ensure you have the instruments to measure
the multiple causes across the age span.



Assess trauma exposure and symptoms of posttraumatic stress

The **Detailed Assessment of Posttraumatic Stress™ (DAPS™)** is a self-report instrument for adults ages 18 to 91 years that assesses peri- and posttraumatic symptoms and associated features related to a specific traumatic event. It provides a detailed assessment of PTSD with an administration time of 30 minutes. Available in paper and pencil, **online via PARiConnect**, and software.


Learn more at parinc.com/DAPS

Evaluate the effects of trauma in adults

The **Trauma Symptom Inventory™-2 (TSI-2™)** is a broadband measure designed to evaluate posttraumatic stress and other psychological sequelae of traumatic events in adults ages 18 to 88 years, including effects of combat, physical abuse, and childhood abuse or neglect. Administration time is 20 minutes. Available in paper and pencil, **online via PARiConnect**, and software. The **TSI-2-A** is an alternate form that does not contain any sexual symptom items.

Learn more at parinc.com/TSI2





Quickly evaluate effects of trauma in children


The **Trauma Symptom Checklist for Children™ (TSCC™)** allows you to measure stress and related psychological symptomatology in children ages 8 to 16 years who have experienced traumatic events. Administration time is 20 minutes. Available in paper and pencil, **online via PARiConnect**, and software. The **TSCC Screening Form (TSCC™-SF)** is also available, takes just 5 minutes to administer, and is available in paper and pencil or **via PARiConnect**. The **TSCC-A** is an alternate form that makes no reference to sexual issues.

Learn more at parinc.com/TSCC or parinc.com/TSCC_SF

The first broadband trauma measure for young children

The **Trauma Symptom Checklist for Young Children™ (TSCYC™)** is the first fully standardized and normed broadband trauma measure for children ages 3 to 12 years who have been exposed to traumatic events such as child abuse, peer assault, and community violence. Administration time is 20 minutes. Available in paper and pencil, **online via PARiConnect**, and software. The **TSCYC Screening Form (TSCYC™-SF)** is also available, takes just 5 minutes to administer, and is available in paper and pencil or **via PARiConnect**.

Learn more at parinc.com/TSCYC and parinc.com/TSCYC_SF



PUT OUR EXPERIENCE TO WORK FOR YOU!

Learn more from
your **PAR Healthcare**
Clinical Assessment Advisor



Melissa Milanak, PhD

813.955.2927 or
mmilanak@parinc.com

Learn more at parinc.com